

Effects of Exercise on Cardiovascular Endurance and the Relationship Between Nutritional Habits and Sports Performance

¹Ayodhya Prasad Pradhan, ²Kamla Sidar, ³Keshav Patel

¹Govt. Vijay Bhushan Singh Dev Girls' College Jashpur (C.G.)

²Government College Barpali, Korba (C.G.)

³Shahid Durvasa Nishad Govt. College Arjunda, Balod (C.G.)

Abstract

Cardiovascular endurance and nutritional habits are important factors influencing physical fitness and sports performance. Regular exercise improves the ability of the cardiovascular and respiratory systems to supply oxygen to working muscles, while appropriate nutrition supports energy production, recovery, and sustained physical activity. This review examines the effects of regular exercise on cardiovascular endurance and explores the relationship between nutritional habits and sports performance, particularly among college-level athletes and physically active young adults. Published research on exercise training, sports nutrition, hydration, and energy availability is examined to understand how these factors contribute to athletic performance. Existing evidence indicates that structured aerobic training can improve cardiorespiratory fitness and endurance capacity. Similarly, adequate carbohydrate intake, protein consumption, hydration, and sufficient energy availability support exercise performance and recovery. Poor dietary practices and inadequate energy intake may reduce training quality, delay recovery, and negatively affect physical performance. Exercise and nutrition are therefore closely related components of athletic development. Their combined influence depends on individual fitness, training intensity, dietary adequacy, and recovery practices. The review highlights the importance of integrating regular exercise with balanced nutritional habits to promote cardiovascular endurance and sports performance, particularly within college sports programmes in India.

Keywords: Physical Exercise, Cardiovascular Endurance, Nutritional Habits, Sports Performance, Aerobic Training, Sports Nutrition, College Athletes.

1. Introduction

Physical fitness plays an important role in maintaining health and achieving satisfactory sports performance. Among the various components of physical fitness, cardiovascular endurance is particularly important because it determines the ability to sustain physical activity over an extended period. Athletes participating in running, cycling, swimming, football, hockey, and other physically demanding sports require sufficient cardiovascular capacity to maintain performance and resist fatigue.

Cardiovascular endurance refers to the ability of the heart, lungs, blood vessels, and skeletal muscles to support prolonged physical activity. Regular aerobic exercise improves the

efficiency of oxygen transport and utilization, helping individuals perform sustained activities with greater efficiency. The American College of Sports Medicine emphasizes that appropriate exercise frequency, intensity, duration, and progression contribute to the development and maintenance of cardiorespiratory fitness [1].

Although physical training is essential for improving endurance, exercise alone does not determine sports performance. Nutritional habits influence the availability of energy, recovery after physical activity, and the body's ability to adapt to training. Athletes require adequate nutrients to support muscular activity, maintain health, and perform consistently during practice and competition.

Thomas et al. [2] established that well-planned nutritional strategies can improve athletic performance and recovery. Dietary requirements vary according to the type of sport, exercise duration, training intensity, and individual characteristics. Carbohydrates provide an important energy source during endurance activities, while proteins support muscle repair and adaptation. Dietary fats, vitamins, minerals, and fluids also contribute to normal physiological functioning.

In India, college sports programmes provide opportunities for students to participate in competitive and recreational physical activities. However, nutritional requirements may not always receive the same attention as physical training. Academic schedules, food preferences, financial limitations, and the availability of balanced meals can influence daily dietary practices.

Understanding the combined importance of exercise and nutrition is therefore necessary for developing effective sports training programmes. Regular exercise can improve cardiovascular endurance, but inadequate nutrition may affect training quality and recovery. Similarly, balanced nutrition supports physical performance but cannot replace the physiological adaptations produced by consistent exercise.

This review examines the effects of regular exercise on cardiovascular endurance and discusses the relationship between nutritional habits and sports performance. Particular attention is given to the implications for college athletes and physical education programmes in India.

2. Review Methodology

A narrative review approach was adopted to examine published evidence relating to exercise, cardiovascular endurance, nutritional habits, and sports performance. Relevant peer-reviewed research articles, scientific reviews, and professional consensus statements were considered.

The review focused on three major areas: the physiological effects of exercise on cardiovascular endurance, the contribution of nutrition to athletic performance, and the combined role of training and dietary practices in endurance development.

Evidence concerning aerobic training, exercise prescription, carbohydrate availability, protein intake, hydration, and energy deficiency was examined. Studies involving athletes and

physically active adults were included because their findings provide a scientific basis for understanding sports performance.

Greater attention was given to evidence relevant to college-level athletes and practical applications within Indian educational institutions. Findings were organized according to major physiological and nutritional factors. The available evidence was interpreted qualitatively, without conducting an independent statistical meta-analysis.

3. Review of Related Literature

Scientific research has established that cardiovascular endurance is influenced by regular physical training and that nutritional strategies contribute to athletic performance and recovery.

Garber et al. [1] presented an American College of Sports Medicine position statement describing the quantity and quality of exercise required to develop and maintain cardiorespiratory, muscular, and neuromotor fitness. The recommendations emphasized the importance of exercise intensity, duration, frequency, and gradual progression. These principles remain important for designing structured endurance training programmes.

Thomas et al. [2] examined nutrition and athletic performance in a joint professional position statement. The authors emphasized that appropriate energy intake, carbohydrate availability, protein consumption, and hydration are important for exercise performance and recovery. They also highlighted the need to adjust nutritional strategies according to training demands and individual requirements.

Burke et al. [3] reviewed the process of muscle glycogen restoration following exercise. The authors explained the importance of carbohydrate availability in replenishing muscle glycogen and supporting recovery between demanding exercise sessions. Their findings are particularly relevant to athletes participating in repeated endurance training or competitions.

Sawka et al. [4] examined exercise and fluid replacement through an American College of Sports Medicine position statement. The authors discussed the effects of dehydration and the importance of appropriate fluid intake during physical activity. They emphasized that hydration requirements depend on environmental conditions, exercise duration, sweat losses, and individual characteristics.

Mountjoy et al. [5] presented the International Olympic Committee's updated consensus statement on Relative Energy Deficiency in Sport. The report explained that prolonged inadequate energy availability can negatively influence health and sports performance. This evidence demonstrates that athletes require sufficient dietary energy to support both training demands and normal physiological functioning.

Maughan et al. [6] examined dietary supplements and athletic performance in an International Olympic Committee consensus statement. Their findings emphasized that basic nutritional adequacy should receive priority over unnecessary supplement use. Although selected

supplements may offer benefits in specific circumstances, routine dietary practices remain the foundation of sports nutrition.

Mujika and Padilla [7] discussed precompetition tapering strategies and explained that reducing accumulated training fatigue before important competitions can support performance. Their findings highlight the importance of recovery and appropriate training management in addition to exercise intensity and volume.

The reviewed literature indicates that exercise and nutrition influence sports performance through closely related processes. Aerobic training develops the physiological capacity for sustained activity, while adequate dietary intake provides the energy and nutrients necessary to support exercise, recovery, and adaptation.

Table 1. Summary of Selected Research Studies

Author and Year	Research Focus	Main Findings
Garber et al. (2011) [1]	Exercise prescription and cardiorespiratory fitness	Structured exercise supports cardiovascular fitness
Thomas et al. (2016) [2]	Nutrition and athletic performance	Appropriate nutrition improves performance and recovery
Burke et al. (2017) [3]	Postexercise glycogen restoration	Carbohydrate availability supports recovery
Sawka et al. (2007) [4]	Hydration and exercise	Fluid replacement is important for maintaining exercise performance
Mountjoy et al. (2023) [5]	Energy deficiency in athletes	Inadequate energy availability may impair health and performance
Maughan et al. (2018) [6]	Dietary supplements and performance	Food-based nutrition remains fundamental
Mujika and Padilla (2003) [7]	Training taper and sports performance	Appropriate recovery and reduced fatigue support competition performance

4. Effects of Exercise on Cardiovascular Endurance

4.1 Development of Cardiovascular Fitness

Cardiovascular endurance depends on the ability of the body to deliver oxygen to active muscles and use it efficiently during prolonged exercise. Regular physical training improves cardiovascular function and supports the physiological adaptations required for sustained activity.

Aerobic activities such as running, cycling, swimming, brisk walking, and continuous sports training increase the demand for oxygen and energy. When performed consistently, these activities can improve aerobic capacity and the ability to sustain physical effort.

One important measure of cardiovascular fitness is maximal oxygen uptake, commonly expressed as $\text{VO}_{2\text{max}}$. This measure represents the maximum rate at which oxygen can be taken up and used during intense physical activity. Higher $\text{VO}_{2\text{max}}$ values are generally associated with greater aerobic capacity, although sports performance also depends on movement economy, exercise tolerance, and sport-specific skills.

Garber et al. [1] emphasized that cardiovascular fitness can be improved through appropriately prescribed aerobic exercise. The magnitude of improvement depends on an individual's initial fitness level, training intensity, frequency, and duration.

4.2 Influence of Exercise Intensity and Duration

Exercise intensity is an important factor determining cardiovascular adaptation. Moderate-intensity exercise can improve general aerobic fitness, while appropriately supervised vigorous training may produce additional adaptations in individuals capable of tolerating higher workloads.

Training duration also influences endurance development. Short exercise sessions may be suitable for beginners, whereas trained athletes often require longer or more demanding sessions to meet performance objectives.

However, increasing training intensity or duration without sufficient recovery may result in accumulated fatigue and reduced training quality. Mujika and Padilla [7] highlighted the importance of managing training load and recovery before major competitions.

For college athletes, endurance programmes should be developed gradually according to physical ability, sporting requirements, and previous training experience. Training progression should be individualized rather than based on a single schedule for all participants.

4.3 Exercise and Resistance to Fatigue

Fatigue during prolonged physical activity may arise from several factors, including changes in muscle energy availability, exercise intensity, heat stress, and neuromuscular demands.

Regular endurance training improves the ability to sustain physical activity and helps athletes tolerate repeated exercise demands. Training adaptations can reduce the relative physiological effort required to perform a given workload.

Nevertheless, resistance to fatigue depends on more than exercise training. Muscle glycogen availability, fluid balance, sleep, and recovery practices influence how effectively athletes maintain performance.

Burke et al. [3] emphasized the importance of restoring muscle glycogen after demanding exercise. This suggests that endurance development requires both regular training and adequate nutritional support.

4.4 Assessment of Cardiovascular Endurance

Cardiovascular endurance can be assessed using laboratory and field-based methods. Laboratory measurements of VO_2max provide a direct assessment of maximal aerobic capacity when appropriate equipment and protocols are available.

Field tests such as the Cooper 12-minute run, multistage shuttle run, and timed distance tests are commonly used in physical education settings. These tests provide practical estimates of endurance capacity, although their accuracy depends on the testing procedure and participant characteristics.

In Indian colleges, field-based endurance tests may be particularly useful where advanced laboratory facilities are unavailable. Standardized testing conditions and appropriate supervision are important for obtaining reliable results.

5. Relationship Between Nutritional Habits and Sports Performance

5.1 Role of Balanced Nutrition

Balanced nutrition is essential for maintaining health, supporting physical activity, and enabling recovery from training. An appropriate diet supplies carbohydrates, proteins, fats, vitamins, minerals, and fluids in quantities that correspond to individual requirements.

Athletes performing regular exercise often have greater energy needs than sedentary individuals. However, nutritional requirements differ according to body size, training load, sport, environmental conditions, and personal health.

Thomas et al. [2] emphasized that nutrition should be planned according to the demands of training and competition. Adequate dietary intake supports exercise performance, while insufficient nutrition may reduce the ability to sustain demanding physical activity.

For college athletes, balanced meals can be prepared using commonly available foods such as rice, wheat products, pulses, vegetables, fruits, milk products, eggs, and other suitable protein sources.

5.2 Carbohydrate Intake and Endurance Performance

Carbohydrates are an important energy source during moderate- and high-intensity exercise. They are stored primarily as glycogen in skeletal muscles and the liver.

During prolonged physical activity, carbohydrate availability contributes to the ability to maintain exercise intensity. When glycogen stores become substantially depleted, athletes may experience difficulty sustaining demanding workloads.

Burke et al. [3] described the importance of carbohydrate consumption in restoring muscle glycogen after exercise. Recovery strategies become particularly important when athletes perform repeated training sessions with limited time between them.

Appropriate carbohydrate intake should therefore be considered in relation to exercise intensity, duration, and recovery needs. Foods such as rice, chapati, potatoes, fruits, and cereals can provide useful carbohydrate sources within a balanced diet.

However, carbohydrate requirements are not identical for all athletes. Individuals participating in light recreational activity generally require different amounts than those completing prolonged endurance training.

5.3 Protein Intake and Muscular Recovery

Protein contributes to muscle maintenance, repair, and adaptation following physical exercise. Repeated training produces physiological demands that require adequate dietary protein and total energy intake.

Protein-rich foods such as milk, curd, pulses, soy products, eggs, fish, and meat can support athletes' nutritional requirements according to their dietary preferences.

Thomas et al. [2] emphasized that protein intake should be considered in relation to training demands and recovery requirements. Although protein is important for muscular adaptation, excessive consumption does not automatically improve endurance performance.

For college athletes, a balanced distribution of protein-rich foods across daily meals may be more practical than depending on expensive supplements. Adequate carbohydrate intake and overall energy availability remain important components of recovery.

5.4 Hydration and Exercise Performance

Hydration is essential for regulating body temperature, supporting circulation, and maintaining normal physiological function during exercise.

Physical activity increases fluid loss through sweating, particularly in hot environments. Inadequate fluid replacement may contribute to dehydration, reduced exercise tolerance, and increased physiological strain.

Sawka et al. [4] emphasized the importance of developing hydration strategies based on exercise conditions and individual sweat losses.

The Indian climate may create additional hydration challenges for athletes training outdoors, especially during hot and humid periods. Appropriate access to drinking water and sensible exercise scheduling are therefore important.

Hydration requirements vary among individuals. Both insufficient fluid intake and excessive fluid consumption can create health concerns, making individualized drinking practices preferable to rigid recommendations.

5.5 Energy Availability and Dietary Deficiencies

Sufficient energy availability is necessary to support physical training, recovery, and normal physiological functions.

Athletes who repeatedly consume less energy than required may experience impaired recovery, reduced training adaptation, and other health concerns. The International Olympic Committee identifies prolonged problematic low energy availability as an important factor associated with Relative Energy Deficiency in Sport [5].

This condition can affect athletes of different sexes and sporting backgrounds. Its consequences may involve metabolic, hormonal, skeletal, and performance-related disturbances.

For college athletes, irregular meals, restrictive dieting, limited food availability, or inadequate nutritional knowledge may contribute to insufficient dietary intake.

Regular meals and appropriate nutritional guidance are important for supporting exercise participation and reducing avoidable nutritional problems.

5.6 Dietary Habits and the Indian College Environment

Nutritional practices among college students may be influenced by academic schedules, hostel facilities, food costs, family dietary patterns, and personal preferences.

Students involved in sports may find it challenging to coordinate meals with morning training, classroom attendance, evening practice, and examinations.

Affordable foods available in India can contribute to a balanced sports diet when selected in appropriate combinations. Rice, roti, dal, seasonal vegetables, fruits, milk, curd, groundnuts, and other locally available foods can provide important nutrients.

Sports nutrition education may help college athletes understand meal planning, hydration, training-day energy requirements, and the limitations of unnecessary supplements.

Maughan et al. [6] emphasized that supplements should not replace a nutritionally adequate diet. Therefore, institutional efforts should focus first on practical food-based strategies.

Table 2. Nutritional Factors and Their Relationship With Sports Performance

Nutritional Factor	Main Function	Relationship With Performance
Carbohydrates	Provide energy and replenish glycogen	Support sustained exercise and recovery
Protein	Supports tissue repair and adaptation	Contributes to recovery and muscle maintenance
Dietary fats	Provide energy and essential fatty acids	Support health and energy requirements

Vitamins and minerals	Support metabolic and physiological functions	Deficiencies may impair health and performance
Water and electrolytes	Support fluid balance and temperature regulation	Help maintain exercise tolerance
Adequate energy intake	Supports training and normal body functions	Reduces risks associated with low energy availability
Regular balanced meals	Support consistent nutrient intake	Help meet daily training and recovery needs

6. Discussion

The available literature indicates that regular exercise and appropriate nutritional habits are closely related to cardiovascular endurance and sports performance. Physical training develops the physiological capacity to perform sustained exercise, whereas nutrition helps provide the energy and nutrients required for that activity.

Aerobic exercise improves cardiorespiratory fitness through repeated physiological demands on the cardiovascular and respiratory systems. However, the benefits of training depend on exercise intensity, duration, progression, and recovery.

Nutrition influences performance through energy availability, muscle glycogen storage, hydration, tissue repair, and maintenance of general health. Inadequate dietary intake may prevent athletes from completing training at the intended intensity or recovering effectively between sessions.

An important observation is that exercise and nutrition do not operate independently. Even a well-designed training programme may be less effective when an athlete experiences persistent energy deficiency or inadequate recovery. Similarly, balanced nutrition alone cannot produce the cardiovascular adaptations associated with regular aerobic training.

The importance of carbohydrate availability is particularly evident during prolonged endurance exercise. Sufficient carbohydrate intake can support sustained workloads and recovery, while protein and other nutrients contribute to adaptation and tissue maintenance.

Hydration also deserves attention in Indian sports environments. Outdoor training under hot conditions may increase fluid losses and physiological strain. Appropriate hydration and exercise scheduling can help athletes manage these demands.

The reviewed evidence supports practical nutrition education within college sports programmes. Students may benefit from understanding balanced meals, meal timing, hydration, food selection, and recovery practices. Such guidance should consider affordability, local food availability, personal preferences, and sporting requirements.

However, several limitations should be recognized. Much of the available evidence comes from controlled studies, professional athletes, or international populations. Findings from these groups cannot always be applied directly to Indian college students with different training backgrounds and resources.

Additionally, sports performance is influenced by many variables beyond nutrition and cardiovascular endurance, including muscular strength, technical ability, motivation, sleep, coaching, and competition conditions. Therefore, the relationship between nutritional habits and sports performance should not be interpreted as a simple cause-and-effect association in every situation.

Further research involving Indian college athletes should assess dietary intake, exercise participation, cardiovascular endurance, and objective performance measures together. Longitudinal and intervention-based studies may help establish how changes in nutrition and training contribute to endurance development over time.

7. Conclusion

Regular exercise and balanced nutritional habits are important components of cardiovascular endurance and sports performance. Aerobic training improves the ability of the cardiovascular and respiratory systems to support prolonged physical activity, while appropriate nutrition contributes to energy availability, recovery, and physiological adaptation.

The reviewed literature indicates that carbohydrates, proteins, adequate energy intake, and hydration play important roles in supporting athletic activity. Poor nutritional practices may contribute to fatigue, inadequate recovery, and reduced training effectiveness.

Exercise and nutrition should therefore be considered complementary elements of sports preparation rather than independent factors. In Indian colleges, structured endurance training combined with practical nutrition education may help students develop healthier routines and improve their readiness for sports participation.

Greater attention to locally available balanced diets, hydration, recovery practices, and individualized exercise programmes can strengthen physical education initiatives. Further research involving Indian college athletes is required to evaluate the combined long-term effects of exercise and nutritional habits on cardiovascular endurance and measurable sports performance.

References

[1] C. E. Garber et al., "American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise," *Medicine & Science in Sports & Exercise*, vol. 43, no. 7, pp. 1334–1359, 2011. doi: 10.1249/MSS.0b013e318213febf.

- [2] D. T. Thomas, K. A. Erdman, and L. M. Burke, "Position of the Academy of Nutrition and Dietetics, Dietitians of Canada, and the American College of Sports Medicine: Nutrition and athletic performance," *Journal of the Academy of Nutrition and Dietetics*, vol. 116, no. 3, pp. 501–528, 2016. doi: 10.1016/j.jand.2015.12.006.
- [3] L. M. Burke, L. J. C. van Loon, and J. A. Hawley, "Postexercise muscle glycogen resynthesis in humans," *Journal of Applied Physiology*, vol. 122, no. 5, pp. 1055–1067, 2017. doi: 10.1152/jappphysiol.00860.2016.
- [4] M. N. Sawka et al., "American College of Sports Medicine position stand. Exercise and fluid replacement," *Medicine & Science in Sports & Exercise*, vol. 39, no. 2, pp. 377–390, 2007. doi: 10.1249/mss.0b013e31802ca597.
- [5] M. Mountjoy et al., "2023 International Olympic Committee's (IOC) consensus statement on Relative Energy Deficiency in Sport (REDs)," *British Journal of Sports Medicine*, vol. 57, no. 17, pp. 1073–1097, 2023. doi: 10.1136/bjsports-2023-106994.
- [6] R. J. Maughan et al., "IOC consensus statement: Dietary supplements and the high-performance athlete," *British Journal of Sports Medicine*, vol. 52, no. 7, pp. 439–455, 2018. doi: 10.1136/bjsports-2018-099027.
- [7] I. Mujika and S. Padilla, "Scientific bases for precompetition tapering strategies," *Medicine & Science in Sports & Exercise*, vol. 35, no. 7, pp. 1182–1187, 2003. doi: 10.1249/01.MSS.0000074448.73931.11.